



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities.

CALIFORNIA—*Los Angeles*.—Month of March, 1898. Estimated population, 103,000. Total deaths, 161, including phthisis pulmonalis, 31; enteric fever, 2; diphtheria, 4, and measles, 1.

Oakland.—Month of March, 1898. Estimated population, 75,000. Total deaths, 103, including phthisis pulmonalis, 19, and diphtheria, 2.

San Diego.—Month of March, 1898. Estimated population, 22,000. Total deaths, 18, including 3 from phthisis pulmonalis.

CONNECTICUT.—Month of March, 1898. Reports to the State board of health from 167 towns, having an aggregate population of 881,192, show a total of 1,272 deaths, including phthisis pulmonalis, 133; enteric fever, 7; scarlet fever, 2; diphtheria, 30; measles, 6, and whooping cough, 10.

ILLINOIS—*Chicago*.—Month of March, 1898. Estimated population, 1,750,000. Total deaths, 2,316, including phthisis pulmonalis, 262; enteric fever, 41; scarlet fever, 10; diphtheria, 43; measles, 7, and whooping cough, 25.

LOUISIANA—*Baton Rouge*.—Month of October, 1897. Estimated population, white, 5,000; colored, 7,000; total, 12,000. Deaths, white, 4; colored, 10; total, 14, including phthisis pulmonalis, 1, and enteric fever, 1.

Month of November, 1897. Deaths, white, 4; colored, 11; total, 15, including yellow fever, 1, and enteric fever, 1.

Month of December, 1897. Deaths, white, 8; colored, 14; total, 22, including 1 from enteric fever.

New Orleans.—Month of October, 1897. Estimated population, white, 195,000; colored, 80,000; total, 275,000. Deaths, white, 451; colored, 199; total, 650, including yellow fever, 158; enteric fever, 11; diphtheria, 3; phthisis pulmonalis, 58, and whooping cough, 1.

Month of November, 1897. Deaths, white, 409; colored, 171; total, 580, including yellow fever, 105; enteric fever, 19; phthisis pulmonalis, 70, and diphtheria, 1.

Month of December, 1897. Deaths, white, 347; colored, 219; total, 566, including yellow fever, 7; enteric fever, 9; and phthisis pulmonalis, 78.

Shreveport.—Month of October, 1897. Estimated population, white, 7,500; colored, 7,500; total, 15,000. Deaths, white, 10; colored, 26; total, 36, including phthisis pulmonalis, 5; enteric fever, 1, and diphtheria, 1.

Month of November, 1897. Deaths, white, 11; colored, 17; total, 28, including 4 from phthisis pulmonalis.

Month of December, 1897. Deaths, white, 8; colored, 31; total, 39, including phthisis pulmonalis, 6, and croup, 1.

MICHIGAN.—Month of March, 1898. Reports to the Secretary of State of the State of Michigan from 83 counties, having an aggregate

population of 2,155,421, show a total of 2,686 deaths, including phthisis pulmonalis, 233; enteric fever, 23; scarlet fever, 13; diphtheria, 29; measles, 18, and whooping cough, 13. Week ended April 9, 1898. Reports to the State board of health from 61 observers indicate that pneumonia increased and intermittent fever decreased in area of prevalence. Phthisis pulmonalis was reported present at 136, measles at 55, scarlet fever at 26, enteric fever at 23, diphtheria at 22, whooping cough at 8, and smallpox at 1 place (Ionia).

MINNESOTA—*Minneapolis*.—Month of March, 1898. Estimated population, 225,602. Total deaths, 219, including phthisis pulmonalis, 29; enteric fever, 8; diphtheria, 1; measles, 29, and whooping cough, 1.

MISSOURI—*Kansas City*.—Month of January, 1898. Estimated population, 200,000. Total deaths, 195, including phthisis pulmonalis, 28; enteric fever, 6; scarlet fever, 2, and diphtheria, 7.

NEW HAMPSHIRE—*Concord*.—Month of March, 1898. Estimated population, 18,000. Total deaths, 29, including 1 from phthisis pulmonalis.

NEW YORK—*Auburn*.—Month of March, 1898. Estimated population, 30,000. Total deaths, 36, including 7 from phthisis pulmonalis.

Buffalo.—Month of March, 1898. Estimated population, 360,000. Total deaths, 327, including phthisis pulmonalis, 36; enteric fever, 11; diphtheria, 7; scarlet fever, 3, and whooping cough, 1.

OHIO—*Cleveland*.—Month of March, 1898. Estimated population, 370,000. Total deaths, 491, including phthisis pulmonalis, 54; enteric fever, 3; scarlet fever, 1; measles, 6; diphtheria, 9, and whooping cough, 1.

Columbus.—Month of March, 1898. Estimated population, 135,000. Total deaths, 153, including phthisis pulmonalis, 32; enteric fever, 1; scarlet fever, 1; diphtheria, 1; measles, 4, and whooping cough, 1.

PENNSYLVANIA—*Williamsport*.—Month of March, 1898. Estimated population, 35,000. Total deaths, 26. No deaths from contagious diseases.

TENNESSEE—*Memphis*.—Month of March, 1898. Estimated population, white, 31,430; colored, 30,317; total, 61,747. Deaths, white, 55; colored, 71; total, 126, including phthisis pulmonalis, 18, and enteric fever, 1.

WASHINGTON—*Seattle*.—Month of March, 1898. Estimated population, 70,000. Total deaths, 78, including phthisis pulmonalis, 7; enteric fever, 2; diphtheria, 6; measles, 3, and whooping cough, 1.